

# Structure of Extreme Correlated Equilibria

Noah D. Stein, Asuman Ozdaglar, and Pablo A. Parrilo<sup>\*†</sup>

January 29, 2010

## Abstract

We exhibit the rich structure of the set of correlated equilibria by analyzing the simplest of polynomial games: the mixed extension of matching pennies. We show that while the correlated equilibrium set is convex and compact, the structure of its extreme points can be quite complicated. In finite games there can be a superexponential separation between the number of extreme Nash and extreme correlated equilibria. In polynomial games there can exist extreme correlated equilibria which are not finitely supported; we construct a large family of examples using techniques from ergodic theory. We show that in general the set of correlated equilibrium distributions of a polynomial game cannot be described by conditions on finitely many joint moments, in marked contrast to the set of Nash equilibria which is always expressible in terms of finitely many moments.

## 1 Introduction

Correlated equilibria are a natural generalization of Nash equilibria introduced by Aumann [1]. They are defined to be joint probability distributions over the players' strategy spaces, such that if each player receives a private recommendation sampled according to the distribution, no player has an incentive to deviate unilaterally from playing his recommended strategy. In finite games the set of correlated equilibria is a compact convex polytope, and therefore seemingly much simpler than the set of Nash equilibria, which can be essentially any algebraic variety [4]. Even in the simple case of two-player finite games, the set of Nash equilibria is a union of finitely many polytopes: seemingly more complicated than the set of correlated equilibria.

Nonetheless we will see that there are two-player zero-sum games in which the set of correlated equilibria has many more extreme points than the set of Nash equilibria has. This behavior does not seem to be pathological in any way: it occurs in very simple finite

---

<sup>\*</sup>Department of Electrical Engineering, Massachusetts Institute of Technology: Cambridge, MA 02139. [nstein@mit.edu](mailto:nstein@mit.edu), [asuman@mit.edu](mailto:asuman@mit.edu), and [parrilo@mit.edu](mailto:parrilo@mit.edu).

<sup>†</sup>This research was funded in part by National Science Foundation grants DMI-0545910 and ECCS-0621922 and AFOSR MURI subaward 2003-07688-1.

games and the simplest of infinite games. We take this as evidence that this complexity is likely to be quite common.

## Contributions

- We give a family of examples of two-player zero-sum finite games in which the set of Nash equilibria has polynomially many extreme points (Section 3), while the set of correlated equilibria has factorially many extreme points (Section 4).
- We give a related example of a continuous game with strategy sets equal to  $[-1, 1]$  and bilinear utility functions. This game is just the mixed extension of matching pennies, but we show that it has extreme correlated equilibria with arbitrarily large finite support (Proposition 4.5) and also with infinite support (Proposition 4.12). This is in contrast to the extreme Nash equilibria, which always have uniformly bounded finite support in zero-sum games with polynomial utilities [12].
- Comparing Proposition 4.14 with this example shows that in general there is no finite-dimensional description of the set of correlated equilibria of a zero-sum polynomial game. Such a description for the Nash equilibria has been known for over fifty years [12].

The examples are closely related – the finite game examples are just restrictions of the strategy spaces in the infinite game example to fixed finite sets. This allows us to analyze both examples on equal footing.

**Related Literature** The geometry of Nash and correlated equilibria has been studied extensively. Therefore we only mention work below if it is directly connected to ours and we do not attempt to be exhaustive.

The result most closely related to the present paper states that in two-player finite games, extreme Nash equilibria are a subset of the extreme correlated equilibria. Cripps [3] and Evangelista and Raghavan [7] proved this independently. This result shows that it makes sense to compare the number of extreme Nash and correlated equilibria. It also raises the natural question of whether all extreme Nash equilibria could be enumerated efficiently by enumerating the extreme correlated equilibria. We show that there can be many more extreme correlated equilibria than extreme Nash equilibria, answering this question in the negative.

In a similar vein, Nau et al. [17] show that for non-trivial finite games with any number of players, the Nash equilibria lie on the boundary of the correlated equilibrium polytope. With three or more players, the Nash equilibria need not be extreme correlated equilibria. For example consider the three-player poker game analyzed by Nash in [16] which has rational payoffs, hence rational extreme correlated equilibria, but whose unique Nash equilibrium uses irrational probabilities.

Separable games, a generalization of polynomial games, were first studied around the 1950's by Dresher, Karlin, and Shapley in papers such as [6], [5], and [13], which were later

combined in Karlin’s book [12]. Their work focuses on the zero-sum case, which contains some of the key ideas for the nonzero-sum case. In particular, they show how to replace the infinite-dimensional mixed strategy spaces with finite-dimensional moment spaces. Carathéodory’s theorem [2] then applies to show that finitely-supported Nash equilibria exist.

There are many similarities between separable games and finite games whose payoff matrices satisfy low-rank conditions. Lipton et al. [14] consider two-player finite games and provide bounds on the cardinality of the support of extreme Nash equilibrium strategies in terms of the ranks of the payoff matrices. The main technical tool here is again Carathéodory’s theorem.

Germano and Lugosi show that in finite games with three or more players there exist correlated equilibria with smaller support than one might expect for Nash equilibria [9]. The proof is geometrical; it essentially views correlated equilibria as living in a subspace of low codimension and it too uses Carathéodory’s theorem [2].

The bounds on the support of equilibria in finite and separable games of the previous three paragraphs are all synthesized in [21]; the portion on Nash equilibria has appeared in [22]. To produce upper bounds on the minimal support of correlated equilibria which depend only on the rank of the payoff matrices and not on the size of the strategy sets, this work does not bound the support of all extreme correlated equilibria, but rather only those whose support is contained inside a Nash equilibrium of small support, which must exist. Similar results hold for polynomial games with, for example, degree used in place of rank (the notions of degree and rank are generalized in [21] and [22]).

This work left open the question of whether all extreme correlated equilibria have support size which can be bounded in terms of the rank of the payoff matrices, independently of the size of the strategy sets. Here we show that this is not the case, because our examples have payoffs which are of rank 1 and extreme correlated equilibria of arbitrarily large, even infinite, support.

Correlated equilibria without finite support have been defined and studied by several authors. An important example of this line of research is the paper by Hart and Schmeidler [11]. The definition of correlated equilibria presented in [11] is convenient for proving some theoretical results (they focus on existence) but not usually for computation.

The authors of the present paper have developed several equivalent characterizations of correlated equilibria in continuous games which are more suitable for computation [23]. One of these forms the basis for the analysis in Section 4 below. Other such characterizations lead to algorithms for approximating correlated equilibria of continuous games [23].

**Outline** The remainder of this paper is organized as follows. Section 2 introduces the examples to be studied. In Section 3 we define and compute the extreme Nash equilibria of these examples, counting them in the finite game example. Then we define and analyze the extreme correlated equilibria in Section 4. This analysis is somewhat long and at times technical, so we present a detailed roadmap before beginning. We close with Section 5, where we outline directions for future work.

$(u_X, u_Y)$	$x = -1$	$x = 1$
$y = -1$	$(1, -1)$	$(-1, 1)$
$y = 1$	$(-1, 1)$	$(1, -1)$

Table 1: Utilities for matching pennies

## 2 Description of the examples

First we fix notation. Given a topological space  $S$ , the set  $\Delta(S)$  will denote the set of Borel probability measures on  $S$  and the set  $\Delta^*(S)$  will denote the set of finite Borel measures on  $S$ . In particular  $\Delta(S)$  is the set of measures in  $\Delta^*(S)$  with unit mass. If  $S$  is finite then  $\Delta(S)$  is a simplex and  $\Delta^*(S)$  is an orthant in  $\mathbb{R}^{|S|}$ . We abuse notation and write the measure of a singleton  $\{p\}$  as  $\mu(p)$  rather than  $\mu(\{p\})$ . For any  $p \in S$ , define  $\delta_p \in \Delta(S)$  to be the measure which assigns unit mass to the point  $p$ . Let  $I = [-1, 1] \subset \mathbb{R}$ .

We will focus mainly on two related examples, one with finite strategy sets and one with infinite strategy sets. We will develop them in parallel by analyzing arbitrary games satisfying the following condition.

**Assumption 2.1.** The game is a zero-sum normal form game with two players, called  $X$  and  $Y$ . The strategy sets  $C_X$  and  $C_Y$  are compact subsets of  $I = [-1, 1]$ , each of which contains at least one positive element and at least one negative element. Player  $X$  chooses a strategy  $x \in C_X$  and player  $Y$  chooses  $y \in C_Y$ . The utility functions are  $u_X(x, y) = xy = -u_Y(x, y)$ , so the game is zero sum<sup>1</sup>.

*Example 2.2.* Fix an integer  $n > 0$ . Let  $C_X$  and  $C_Y$  each have  $2n$  elements,  $n$  of which are positive and  $n$  of which are negative. If we take  $n = 1$  and  $C_X = C_Y = \{-1, 1\}$  then we recover the matching pennies game, as shown in Table 1.

*Example 2.3.* Let  $C_X = C_Y = [-1, 1]$ . Then the game is essentially the mixed extension of matching pennies. That is to say, suppose two players play matching pennies and choose their strategies independently, playing 1 with probabilities  $p \in [0, 1]$  and  $q \in [0, 1]$ . Define the utilities for the mixed extension to be the expected utilities under this random choice of strategies. Letting  $x = 2p - 1$  and  $y = 2q - 1$ , the utility to the first player is  $xy$  and the utility to the second player is  $-xy$ . Therefore this example is the mixed extension of matching pennies, up to an affine scaling of the strategies.

## 3 Extreme Nash equilibria

We will now characterize the extreme points of the sets of Nash equilibria in games satisfying Assumption 2.1. We will fully characterize these and then count them. Since the games are

---

<sup>1</sup>By inspection of the utilities we can see that for any  $C_X$  and  $C_Y$  with at least two points, the rank of this game in the sense of [22] is  $(1, 1)$  (and in fact also in the stronger sense of Theorem 3.3 of that paper). The notion of the rank of a game is related to the rank of the payoff matrices and will not play a significant role in this paper; we merely wish to note that under this definition of complexity of payoffs, the games we consider are extremely simple.

zero-sum, the set of Nash equilibria can be viewed as a Cartesian product of two (weak\*) compact convex sets, the sets of maximin and minimax strategies [10]. The Krein-Milman theorem completely characterizes such sets by their extreme points [19], explaining our focus on extreme points throughout.

We define Nash equilibria in two-player games, which will be sufficient for our purposes, as well as the standard notions of extreme point and extreme ray from convex analysis.

**Definition 3.1.** A **Nash equilibrium** is a pair  $(\sigma, \tau) \in \Delta(C_X) \times \Delta(C_Y)$  such that  $u_X(x, \tau) \leq u_X(\sigma, \tau)$  for all  $x \in C_X$  and  $u_Y(\sigma, y) \leq u_Y(\sigma, \tau)$  for all  $y \in C_Y$  (where we extend utilities by expectation in the usual fashion  $u_X(x, \tau) = \int u_X(x, y) d\tau(y)$ , etc.).

In other words, a Nash equilibrium is a strategy pair in which each player is playing a best reply to his opponent's strategy.

**Definition 3.2.** A point  $x$  in a (usually convex) subset  $K$  of a real vector space is an **extreme point** if  $x = \lambda y + (1 - \lambda)z$  for  $y, z \in K$  and  $\lambda \in (0, 1)$  implies  $x = y = z$ .

The related notion of extreme ray will not be used until the next section, but we record it here for comparison.

**Definition 3.3.** A convex set  $K$  such that  $x \in K$  and  $\lambda \geq 0$  implies  $\lambda x \in K$  is called a **convex cone**. A point  $x \neq 0$  is an **extreme ray** of the convex cone  $K$  if  $x = y + z$  and  $y, z \in K$  implies that  $y$  or  $z$  is a scalar multiple of  $x$ .

The Nash equilibria of games satisfying Assumption 2.1 take the following particularly simple form.

**Proposition 3.4.** *A pair  $(\sigma, \tau) \in \Delta(C_X) \times \Delta(C_Y)$  is a Nash equilibrium of a game satisfying Assumption 2.1 if and only if  $\int x d\sigma(x) = \int y d\tau(y) = 0$ .*

*Proof.* If  $\int x d\sigma(x) = 0$  then  $u_Y(\sigma, y) = 0$  for all  $y \in C_Y$ , so any  $\tau \in \Delta(C_Y)$  is a best response to  $\sigma$ . If  $\int y d\tau(y) = 0$  as well then  $\sigma$  is also a best response to  $\tau$ , so  $(\sigma, \tau)$  is a Nash equilibrium.

Suppose for a contradiction that there exists a Nash equilibrium  $(\sigma, \tau)$  such that  $\int x d\sigma(x) > 0$ ; the other cases are similar. Player  $y$  must play a best response, so  $\int y d\tau(y) < 0$ , which is possible by assumption. Player  $x$  plays a best response to that, so  $\int x d\sigma(x) < 0$ , a contradiction.  $\square$

We introduce the notion of extreme Nash equilibrium in two-player zero-sum games. For an extension of this definition to two-player finite games and a proof that extreme Nash equilibria are always extreme points of the set of correlated equilibria in this setting, see [3] or [7].

**Definition 3.5.** In a two-player zero-sum game, **maximin** and **minimax** strategies are those mixed strategies for player  $X$  and  $Y$ , respectively, which appear in a Nash equilibrium. A Nash equilibrium of a zero-sum game is called **extreme** if  $\sigma$  and  $\tau$  are extreme points of the maximin and minimax sets, respectively.

Applying Proposition 3.4 to this definition, we can characterize the extreme Nash equilibria of games satisfying Assumption 2.1.

**Proposition 3.6.** *Consider a game satisfying Assumption 2.1. A pair  $(\sigma, \tau) \in \Delta(C_X) \times \Delta(C_Y)$  is an extreme Nash equilibrium if and only if  $\sigma$  and  $\tau$  are each either  $\delta_0$  or of the form  $\alpha\delta_u + \beta\delta_v$  where  $u < 0$ ,  $v, \alpha, \beta > 0$ ,  $\alpha + \beta = 1$ , and  $\alpha u + \beta v = 0$ .*

*Proof.* By Proposition 3.4 we must show that these distributions are the extreme points of the set of probability distributions having zero mean. Since  $\delta_0$  is an extreme point of the set of probability distributions, it must be an extreme point of the subset which has zero mean. To see that  $\alpha\delta_u + \beta\delta_v$  is also an extreme point, suppose we could write it as a convex combination of two other probability distributions with zero mean. The condition that both be positive measures implies that both must be of the form  $\alpha'\delta_u + \beta'\delta_v$ . But  $\alpha$  and  $\beta$  as specified above are the unique coefficients which make this be a probability measure with zero mean. Therefore  $\alpha' = \alpha$  and  $\beta' = \beta$ , so  $\alpha\delta_u + \beta\delta_v$  cannot be written as a nontrivial convex combination of probability distributions with zero mean, i.e., it is an extreme point.

Suppose  $\sigma$  were an extreme point which was not of one of these types. Then  $\sigma$  could not be supported on one or two points, so either  $[0, 1]$  or  $[-1, 0)$  could be partitioned into two sets of positive measure. We will only treat the first case; the second is similar. Let  $[0, 1] = A \cup B$  where  $A \cap B = \emptyset$  and  $\sigma(A), \sigma(B) > 0$ . Since  $\sigma$  has zero mean we must have  $\sigma([-1, 0)) > 0$  as well.

For a set  $D$  we define the restriction measure  $\sigma|_D$  by  $\sigma|_D(C) = \sigma(D \cap C)$  for all  $C$ . Then  $\sigma = \sigma|_A + \sigma|_B + \sigma|_{[-1, 0)}$ . Let  $a = \int_A x d\sigma(x)$ ,  $b = \int_B x d\sigma(x)$ , and  $c = \int_{[-1, 0)} x d\sigma(x)$ . Since  $\sigma([-1, 0)) > 0$  and  $x$  is less than zero everywhere on  $[-1, 0)$ , we must have  $c < 0$ . By assumption  $a + b + c = 0$ . Therefore we can write:

$$\sigma = \left( \sigma|_A + \frac{a}{|c|} \sigma|_{[-1, 0)} \right) + \left( \sigma|_B + \frac{b}{|c|} \sigma|_{[-1, 0)} \right)$$

Being an extreme point of the set of probability measures with zero mean,  $\sigma$  must be an extreme ray of the set of positive measures with first moment equal to zero. But this means that we cannot write  $\sigma = \sigma_1 + \sigma_2$  where the  $\sigma_i$  are positive measures with zero first moment unless  $\sigma_i$  is a multiple of  $\sigma$ . Neither of the measures in parentheses above is a multiple of  $\sigma$ , so we have a contradiction.  $\square$

We illustrate this proposition on both examples introduced in Section 2.

*Example 2.2 (cont'd).* In this case neither  $C_X$  nor  $C_Y$  contains zero, so the only extreme Nash equilibria are those in which  $\sigma$  and  $\tau$  are of the form  $\alpha\delta_u + \beta\delta_v$  for  $u < 0$  and  $v > 0$ . For any choice of  $u$  and  $v$  there are unique  $\alpha$  and  $\beta$  satisfying the conditions of Proposition 3.6. There are  $n$  possible choices for each of  $u$  and  $v$  for each of the two players, so there are  $n^4$  extreme Nash equilibria.

*Example 2.3 (cont'd).* Since  $C_X = C_Y = [-1, 1]$ , there are infinitely many extreme Nash equilibria in this case. However, they are all finitely supported and the size of their support is always either one or two. Furthermore the condition that  $(\sigma, \tau)$  be a Nash equilibrium

is equivalent to both having zero mean. This illustrates the general facts that in games with polynomial utility functions the Nash equilibrium conditions only involve finitely many moments of  $\sigma$  and  $\tau$  and the extreme Nash equilibria (when defined, say for zero-sum games) have uniformly bounded support [12].

## 4 Extreme correlated equilibria

In this section we will show that even in finite games, the number of extreme correlated equilibria can be much larger than the number of extreme Nash equilibria. It makes sense to compare these because all extreme Nash equilibria of a two-player game are automatically extreme correlated equilibria [3, 7].

In the case of polynomial games we will show that there can be extreme correlated equilibria with arbitrarily large finite support and with infinite support. This implies that the set of correlated equilibria cannot be characterized in terms of finitely many joint moments.

**Roadmap** The analysis proceeds in several steps which will be technical at times, so we start with an outline of what follows.

- We begin by defining correlated equilibria in games satisfying Assumption 2.1 using a characterization from [23].
- Proposition 4.4 shows that this characterization can be simplified because of our choice of utility functions.
- We use this characterization to construct a family of finitely supported extreme correlated equilibria in Proposition 4.5.
- Then we note that all extreme correlated equilibria of the games in Example 2.2 are of this form, so this allows us to count the extreme correlated equilibria and determine their asymptotic rate of growth as the number of pure strategies grows.
- Next we introduce some ideas from ergodic theory. With these tools in hand, we construct in Proposition 4.12 a large family of extreme correlated equilibria without finite support for the game in Example 2.3.
- Finally we show that if a set can be represented by finitely many moments then all its extreme points have uniformly bounded finite support. This shows that the set of correlated equilibria of the game in Example 2.3 cannot be represented by finitely many moments and completes the analysis.

We now begin the analysis. Correlated equilibria are meant to capture the notion of a joint distribution of private recommendations to the two players such that neither player can expect to improve his payoff by deviating unilaterally from his recommendation. For finitely supported probability distributions and games satisfying Assumption 2.1, this can be written as per the standard definition (see [15] or [8]):

**Definition 4.1.** A finitely supported probability distribution  $\mu \in \Delta(C_X \times C_Y)$  is a **correlated equilibrium** of a game satisfying Assumption 2.1 if

$$\sum_{y \in C_Y} \mu(x, y)[xy - x'y] \geq 0$$

for all  $x, x' \in C_X$  and

$$\sum_{x \in C_X} \mu(x, y)[xy' - xy] \geq 0$$

for all  $y, y' \in C_Y$  (note that these sums are finite by the assumption on  $\mu$ ).

The standard definition extending this notion to arbitrary (not necessarily finitely supported) distributions is given in [11]. This definition is difficult to compute with, so we will use the following equivalent characterization.

**Theorem 4.2** ([23]). *A probability distribution  $\mu \in \Delta(C_X \times C_Y)$  is a correlated equilibrium of a game satisfying Assumption 2.1 if and only if*

$$\int_{A \times I} (x - x')y d\mu(x, y) \geq 0 \quad \text{and} \quad \int_{I \times A} x(y - y') d\mu(x, y) \leq 0$$

for all  $x' \in C_X$ ,  $y' \in C_Y$ , and measurable  $A \subseteq I$ .

*Proof.* When  $\mu$  is finitely supported this is clearly equivalent to Definition 4.1. The general case is part (1) of Corollary 2.14 in [23] with the present utilities substituted in.  $\square$

Note that these conditions are homogeneous (that is, invariant under positive scaling) in  $\mu$ . The only condition on  $\mu$  that is not homogeneous is the probability measure condition  $\mu(I \times I) = 1$ . We will often ignore this condition for simplicity, referring to a measure  $\mu \in \Delta^*(C_X \times C_Y)$  satisfying the conditions of the theorem as a correlated equilibrium.

**Definition 4.3.** When we need to distinguish these notions, we will refer to a measure  $\mu \in \Delta^*(C_X \times C_Y)$  satisfying the conditions of Theorem 4.2 as a **homogeneous correlated equilibrium** and a measure  $\mu \in \Delta(C_X \times C_Y)$  satisfying the conditions as a **proper correlated equilibrium**. In the context of homogeneous correlated equilibria the term **extreme** will refer to extreme rays; for proper correlated equilibria it will refer to extreme points.

When  $\mu \neq 0$  is a homogenous correlated equilibrium,  $\frac{1}{\mu(I \times I)}\mu$  is a proper correlated equilibrium. The set of homogenous correlated equilibria is a convex cone. The extreme rays of this cone are exactly those measures which are positive multiples of the extreme points of the set of proper correlated equilibria.

The following proposition characterizes correlated equilibria of games satisfying Assumption 2.1 and is analogous to Proposition 3.4 for Nash equilibria. Note how the Nash equilibrium measures were characterized in terms of their moments but the correlated equilibria are not.



**Proposition 4.4.** *For a game satisfying Assumption 2.1 and a measure  $\mu \in \Delta^*(C_X \times C_Y)$ , the following are equivalent:*

1.  $\mu$  is a correlated equilibrium;

2.

$$\kappa_x(A) := \int_{A \times I} xy d\mu(x, y) \quad \text{and} \quad \kappa_y(A) := \int_{I \times A} xy d\mu(x, y)$$

are both the zero measure, i.e., equal zero for all measurable  $A \subseteq I$ ;

3.

$$\lambda_x(A) := \int_{A \times I} y d\mu(x, y) \quad \text{and} \quad \lambda_y(A) := \int_{I \times A} x d\mu(x, y)$$

are both the zero measure.

*Proof.* (1  $\Rightarrow$  2) We will consider only  $\kappa_x$ ;  $\kappa_y$  is similar. The conditions of Theorem 4.2 with  $A = I$  imply that

$$x' \int_{I \times I} y d\mu(x, y) \leq \int_{I \times I} xy d\mu(x, y) \leq y' \int_{I \times I} x d\mu(x, y)$$

for all  $x' \in C_X, y' \in C_Y$ . By assumption it is possible to choose  $x'$  and  $y'$  either positive or negative, so  $\int_{I \times I} xy d\mu(x, y) = 0$ . Furthermore the same argument with any  $A$  implies that  $\int_{A \times I} xy d\mu(x, y) \geq 0$ . Therefore we have

$$0 = \int_{I \times I} xy d\mu(x, y) = \int_{A \times I} xy d\mu(x, y) + \int_{(I \setminus A) \times I} xy d\mu(x, y) \geq 0 + 0 = 0$$

for all  $A$ , so the inequality must be tight and we get  $\int_{A \times I} xy d\mu(x, y) = 0$  for all  $A$ .

(2  $\Rightarrow$  3) Substituting this equation into Theorem 4.2 yields  $x' \int_{A \times I} y d\mu(x, y) \leq 0$  for all  $x' \in C_X$  and all measurable  $A$ . But we can choose  $x'$  to be positive or negative by assumption, so we must have  $\int_{A \times I} y d\mu(x, y) = 0$  for all measurable  $A \subseteq I$ .

(3  $\Rightarrow$  1) Suppose we know that  $\lambda_x$  is the zero measure. We can approximate  $xy$  on  $I \times I$  by functions of the form  $\sum_{k=1}^r x_i y \chi_{A_i}(x)$ , where the measurable sets  $A_i$  have small diameter and partition  $I$ ,  $x_i \in A_i$ , and  $\chi_{A_i}$  is the indicator function for  $A_i$ . By definition of the Lebesgue integral and the dominated convergence theorem [18] we have

$$\int_{A \times I} xy d\mu(x, y) = \int_A x d\lambda_x(x) = \int_A x d0(x) = 0.$$

Thus  $\int_{A \times I} (x - x')y d\mu(x, y) = 0$  for all  $x' \in C_X$  and all  $A$ . □

**Proposition 4.5.** *Fix a game satisfying Assumption 2.1. Let  $k > 0$  be even and  $x_1, \dots, x_{2k}$  and  $y_1, \dots, y_{2k}$  be such that:*

1.  $x_i \in C_X$  and  $y_i \in C_Y$  are all nonzero;

2. the sequence  $x_1, x_3, \dots, x_{2k-1}$  has distinct elements and alternates in sign;
3. the sequence  $y_1, y_3, \dots, y_{2k-1}$  has distinct elements and alternates in sign;
4.  $x_{2i} = x_{2i-1}$  and  $y_{2i} = y_{2i+1}$  for all  $i$  when subscripts are interpreted  $\pmod{2k}$ .

Then  $\mu = \sum_{i=1}^{2k} \frac{1}{|x_i y_i|} \delta_{(x_i, y_i)}$  is an extreme correlated equilibrium.

*Proof.* To show that  $\mu$  is a correlated equilibrium define  $d\kappa(x, y) = xy d\mu(x, y)$ . Then  $\kappa = \sum_{i=1}^{2k} \text{sign}(x_i) \text{sign}(y_i) \delta_{(x_i, y_i)}$ . Defining the projection  $\kappa_x$  as in Proposition 4.4, we have

$$\begin{aligned} \kappa_x &= \sum_{i=1}^{2k} \text{sign}(x_i) \text{sign}(y_i) \delta_{x_i} = \sum_{i=1}^k \text{sign}(x_{2i}) (\text{sign}(y_{2i}) + \text{sign}(y_{2i-1})) \delta_{x_{2i}} \\ &= \sum_{i=1}^k \text{sign}(x_{2i}) (0) \delta_{x_{2i}} = 0, \end{aligned}$$

because  $x_{2i} = x_{2i-1}$  and  $y_{2i}$  differs in sign from  $y_{2i-1}$  by assumption. The same argument shows that  $\kappa_y = 0$ , so  $\mu$  is a correlated equilibrium.

To see that  $\mu$  is extreme, suppose  $\mu = \mu' + \mu''$  where  $\mu'$  and  $\mu''$  are correlated equilibria. Clearly  $\mu' = \sum_{i=1}^{2k} \alpha_i \delta_{(x_i, y_i)}$  for some  $\alpha_i \geq 0$ . Define  $d\kappa' = xy d\mu'(x, y)$ , so  $\kappa' = \sum_{i=1}^{2k} \alpha_i x_i y_i \delta_{(x_i, y_i)}$ . By assumption

$$\kappa'_x = \sum_{i=1}^k x_{2i} (\alpha_{2i-1} y_{2i-1} + \alpha_{2i} y_{2i}) \delta_{x_{2i}}$$

is the zero measure. Since the  $x_{2i}$  are distinct and nonzero we must have  $\alpha_{2i-1} y_{2i-1} + \alpha_{2i} y_{2i} = 0$  for all  $i$ . Similarly since  $\kappa'_y = 0$  we have  $\alpha_{2i+1} x_{2i+1} + \alpha_{2i} x_{2i} = 0$  for all  $i$  (with subscripts interpreted  $\pmod{2k}$ ).

The  $x_i$  and  $y_i$  are all nonzero, so fixing one  $\alpha_i$  fixes all the others by these equations. That is to say, these equations have a unique solution up to multiplication by a scalar, so  $\mu'$  is a positive scalar multiple of  $\mu$ . But the splitting  $\mu = \mu' + \mu''$  was arbitrary, so  $\mu$  is extreme.  $\square$

An argument along the lines of the proof of Proposition 4.5 shows that any finitely supported correlated equilibrium  $\mu$  whose support does not contain any points with  $x = 0$  or  $y = 0$  can be written as  $\mu = \mu' + \mu''$  where  $\mu'$  is a correlated equilibrium and  $\mu''$  is a correlated equilibrium of the form studied in Proposition 4.5. Therefore  $\mu$  cannot be extreme unless it is of this form.

*Example 2.2 (cont'd).* For some examples of the supports of extreme correlated equilibria of games of this type, see Figures 1 and 2.

To count the number of extreme correlated equilibria of this game we must count the number of essentially different sequences of  $x_i$  and  $y_i$  of the type mentioned in Proposition 4.5. Fix  $k$  and let  $k = 2r$  where  $1 \leq r \leq n$ . Note that cyclically shifting the sequences




Figure 1: The support of an extreme correlated equilibrium. In the notation of Proposition 4.5,  $k = 2$ ,  $x_1 = 0.4$ ,  $x_3 = -0.6$ ,  $y_1 = 0.2$ , and  $y_3 = -0.8$ .




Figure 2: The support of another extreme correlated equilibrium. In the notation of Proposition 4.5,  $k = 4$ ,  $x_1 = 0.4$ ,  $x_3 = -0.4$ ,  $x_5 = 0.6$ ,  $x_7 = -0.6$ ,  $y_1 = 0.6$ ,  $y_3 = -0.4$ ,  $y_5 = 0.4$ , and  $y_7 = -0.6$ .

of  $x_i$ 's and  $y_i$ 's by two does not change  $\mu$ , nor does reversing the sequence. Therefore we can assume without loss of generality that  $x_1, y_1 > 0$ . We then have  $n$  possible choices for  $x_1, y_1, x_3$ , and  $y_3$ ,  $n - 1$  possible choices for  $x_5, x_7, y_5$ , and  $y_7$ , etc., for a total of  $\left(\frac{n!}{(n-r)!}\right)^4$  possible choices of the  $x_i$  and  $y_i$ . These will always be essentially different (i.e., give rise to different  $\mu$ ) unless we cyclically permute the sequences of  $x_i$  and  $y_i$  by some multiple of four, in which case the resulting sequence is essentially the same. The number of such cyclic permutations is  $r$ . Therefore the total number of extreme correlated equilibria is

$$e(n) = \sum_{r=1}^n \frac{1}{r} \left( \frac{n!}{(n-r)!} \right)^4.$$

We will see that  $e(n) = \Theta\left(\frac{1}{n}(n!)^4\right)$ . That is to say,  $e(n)$  is asymptotically upper and lower bounded by a constant times  $\frac{1}{n}(n!)^4$ . The expression  $\frac{1}{n}(n!)^4$  is just the final term in the summation for  $e(n)$ , so the lower bound is clear. Define

$$f(n) = \frac{e(n)}{\frac{1}{n}(n!)^4} = \sum_{s=0}^{n-1} \frac{n}{n-s} \cdot \frac{1}{(s!)^4}.$$

Then  $f(n) \geq 1$  for all  $n$ . We will now show that  $f(n)$  is also bounded above. Intuitively this is not surprising as the terms in the summation for  $f(n)$  die off extremely fast as  $s$  grows.

For all  $1 \leq s < n - 1$  we have that the ratio of term  $s + 1$  in the summation to term  $s$  is:

$$\frac{\frac{n}{n-s-1} \cdot \frac{1}{((s+1)!)^4}}{\frac{n}{n-s} \cdot \frac{1}{(s!)^4}} = \frac{n-s}{n-s-1} \cdot \frac{1}{(s+1)^4} \leq \frac{1}{8},$$

so for  $n > 1$  we can bound the sum by a geometric series:

$$f(n) - 1 = \sum_{s=1}^{n-1} \frac{n}{n-s} \cdot \frac{1}{(s!)^4} \leq \frac{n}{n-1} \sum_{t=0}^{\infty} \frac{1}{8^t} = \frac{8n}{7(n-1)} \leq \frac{16}{7}.$$

Therefore  $1 \leq f(n) \leq \frac{23}{7}$  for all  $n$ , so  $e(n) = \Theta\left(\frac{1}{n}(n!)^4\right)$  as claimed. Comparing this to the results of the previous section in which we saw that the number of extreme Nash equilibria of this game is  $n^4$ , we see that in this case there is a super-exponential separation between the number of extreme Nash and the number of extreme correlated equilibria. This implies, for example, that computing all extreme correlated equilibria is not an efficient method for computing all extreme Nash equilibria, even though all extreme Nash equilibria are extreme correlated equilibria and recognizing whether an extreme correlated equilibrium is an extreme Nash equilibrium is easy. There are simply too many extreme correlated equilibria.

Next we will prove a more abstract version of Proposition 4.5 which includes certain extreme points which are not finitely supported. First we need a brief digression to ergodic theory. The first definition is the standard definition of compatibility between a measure and

a transformation on a space. The second definition expresses one notion of what it means for a transformation to “mix up” a space – in this case that the space cannot be partitioned into two sets of positive measure which do not interact under the transformation. Then we state the main ergodic theorem and a corollary which we will apply to exhibit extreme correlated equilibria of games satisfying Assumption 2.1.

**Definition 4.6.** Given a measure  $\mu \in \Delta^*(S)$  on a space  $S$ , a measurable function  $g : S \rightarrow S$  is called **( $\mu$ -)measure preserving** if  $\mu(g^{-1}(A)) = \mu(A)$  for all measurable  $A \subseteq S$ . Note that if  $g$  is invertible (in the measure theoretic sense that an almost everywhere inverse exists), then this is equivalent to the condition that  $\mu(g(A)) = \mu(A)$  for all  $A$ .

**Definition 4.7.** Given a measure  $\mu \in \Delta^*(S)$ , a  $\mu$ -measure preserving transformation  $g$  is called **ergodic** if  $\mu(A \Delta g^{-1}(A)) = 0$  implies  $\mu(A) = 0$  or  $\mu(A) = \mu(S)$ , where  $A \Delta B$  denotes the symmetric difference  $(A \setminus B) \cup (B \setminus A)$ .

*Example 4.8.* Fix a finite set  $S$  and a permutation  $g : S \rightarrow S$ . Let  $\mu$  be counting measure on  $S$ . Then  $g$  is measure preserving. Furthermore, a set  $T$  satisfies  $\mu(g^{-1}(T) \Delta T) = 0$  if and only if  $g^{-1}(T) = T$  if and only if  $T$  is a union of cycles of  $g$ . Therefore  $g$  is ergodic if and only if it consists of a single cycle.

*Example 4.9.* Fix  $\alpha \in \mathbb{R}$ . Let  $S = [0, 1)$  and let  $\mu$  be Lebesgue measure on  $S$ . Define  $g : S \rightarrow S$  by  $g(x) = (x + \alpha) \bmod 1 = (x + \alpha) - \lfloor x + \alpha \rfloor$ . Then  $g$  is  $\mu$ -measure preserving because Lebesgue measure is translation invariant. It can be shown that  $g$  is ergodic if and only if  $\alpha$  is irrational. For a proof and more examples, see [20].

The following is one of the core theorems of ergodic theory. We will only use it to prove the corollary which follows, so it need not be read in detail. The proof can be found in any text on ergodic theory, e.g. [20].

**Theorem 4.10** (Birkhoff’s ergodic theorem). *Fix a probability measure  $\mu$  and a  $\mu$ -measure preserving transformation  $g$ . Then for any  $f \in \mathcal{L}^1(\mu)$ :*

- $\tilde{f}(x) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f(g^k(x))$  exists  $\mu$ -almost everywhere,
- $\tilde{f} \in \mathcal{L}^1(\mu)$ ,
- $\int \tilde{f} d\mu = \int f d\mu$ ,
- $\tilde{f}(g(x)) = \tilde{f}(x)$   $\mu$ -almost everywhere, and
- if  $g$  is ergodic then  $\tilde{f}(x) = \int f d\mu$   $\mu$ -almost everywhere.

**Corollary 4.11.** *Suppose  $\mu$  and  $\nu$  are probability measures such that  $\nu$  is absolutely continuous with respect to  $\mu$ . If a transformation  $g$  preserves both  $\mu$  and  $\nu$  and  $g$  is ergodic with respect to  $\mu$ , then  $\nu = \mu$ .*

*Proof.* Fix any measurable set  $A$ . Let  $f$  be the indicator function for  $A$ , i.e. the function equal to unity on  $A$  and zero elsewhere. Applying Birkhoff's ergodic theorem to  $f$  and  $\mu$  yields  $\tilde{f}(x) = \mu(A)$   $\mu$ -almost everywhere. Since  $\nu$  is absolutely continuous with respect to  $\mu$ ,  $\tilde{f}(x) = \mu(A)$   $\nu$ -almost everywhere also. If we now apply Birkhoff's ergodic theorem to  $\nu$  we get:

$$\nu(A) = \int f d\nu = \int \tilde{f} d\nu = \int \mu(A) d\nu = \mu(A). \quad \square$$

**Proposition 4.12.** *Fix measures  $\nu_1, \nu_2, \nu_3$ , and  $\nu_4 \in \Delta^*((0, 1])$  and maps  $f_i : (0, 1] \rightarrow (0, 1]$  such that  $\nu_{i+1} = \nu_i \circ f_i^{-1}$  (interpreting subscripts mod 4). The portion of the measure  $\mu$  in the  $i^{\text{th}}$  quadrant of  $I \times I$  will be constructed in terms of  $f_i$  and  $\nu_i$ . Define  $j_i : (0, 1] \rightarrow I \times I$  by  $j_1(x) = (x, f_1(x))$ ,  $j_2(x) = (-f_2(x), x)$ ,  $j_3(x) = (-x, -f_3(x))$ , and  $j_4(x) = (f_4(x), -x)$ . Let  $|\kappa| = \sum_{i=1}^4 \nu_i \circ j_i^{-1}$ . If Assumption 2.1 is satisfied,  $\text{supp}|\kappa| \subseteq C_X \times C_Y$ , and  $\frac{1}{|xy|} \in \mathcal{L}^1(|\kappa|)$  then  $d\mu = \frac{1}{|xy|}d|\kappa|$  is a correlated equilibrium.*

*If in addition  $f_4 \circ f_3 \circ f_2 \circ f_1 : (0, 1] \rightarrow (0, 1]$  is ergodic with respect to  $\nu_1$ , then  $\mu$  is extreme.*

*Proof.* First we must show that  $\mu$  is a correlated equilibrium. It is a finite measure by the assumption  $\frac{1}{|xy|} \in \mathcal{L}^1(|\kappa|)$ . Define  $g : I \times I \rightarrow I \times I$  as follows.

$$g(x, y) = \begin{cases} j_1(x) & \text{if } x > 0, y < 0 \\ j_2(y) & \text{if } x > 0, y > 0 \\ j_3(-x) & \text{if } x < 0, y > 0 \\ j_4(-y) & \text{if } x < 0, y < 0 \\ \text{arbitrary} & \text{otherwise} \end{cases}$$

The function  $g$  is  $|\kappa|$ -measure preserving. To see this fix any measurable set  $B \subseteq (0, 1] \times (0, 1]$ . Let  $A = j_1^{-1}(B)$ . Then  $|\kappa|(B) = |\kappa|(A \times (0, 1]) = \nu_1(A)$  by definition of  $|\kappa|$ . But  $g^{-1}(B) = g^{-1}(A \times (0, 1]) = A \times [-1, 0)$ , so  $|\kappa|(g^{-1}(B)) = |\kappa|(A \times [-1, 0)) = \nu_4(j_4^{-1}(A)) = \nu_1(A)$  by assumption. Therefore  $g$  is measure preserving for subsets of  $(0, 1] \times (0, 1]$ . The arguments for the other quadrants are similar and since  $g$  maps each quadrant into a different quadrant,  $g$  is measure preserving on its entire domain.

Using the terminology of Proposition 4.4,  $d\kappa = \text{sign}(x) \text{sign}(y)d|\kappa|$ . We have seen that  $|\kappa|(A \times (0, 1]) = |\kappa|(A \times [-1, 0))$ , so  $\kappa(A \times (0, 1]) = -\kappa(A \times [-1, 0))$ . Since  $\kappa(A \times \{0\}) = 0$ , we have  $\kappa(A \times I) = 0$ , or equivalently,  $\kappa_x(A) = 0$ . A similar argument implies  $\kappa_x(A) = 0$  if  $A \subseteq [-1, 0)$ . Clearly  $\kappa_x(\{0\}) = 0$  by definition of  $\kappa_x$ , so  $\kappa_x$  is the zero measure. A similar argument shows that  $\kappa_y$  is the zero measure, so  $\mu$  is a correlated equilibrium by Proposition 4.4.

Now we will show via several steps that  $\mu$  is extreme. Write  $\mu = \mu_1 + \mu_2$  where the  $\mu_i$  are nonzero correlated equilibria. Since these are all positive measures, the  $\mu_i$  are absolutely continuous with respect to  $\mu$ . Define  $d|\kappa_i| = |xy|d\mu_i$ .

Next we show that  $g$  is  $|\kappa_i|$ -measure preserving. We will demonstrate this fact for  $B \subseteq (0, 1] \times (0, 1]$ . As above, we define  $A = j_1^{-1}(B)$ . Then  $|\kappa_i|(B) = |\kappa_i|(A \times (0, 1])$  since

$(A \times (0, 1]) \triangle B$  has  $|\kappa|$  measure zero and  $|\kappa_i|$  is absolutely continuous with respect to  $|\kappa|$ . Furthermore,  $|\kappa_i|(g^{-1}(B)) = |\kappa_i|(A \times [-1, 0))$ . But  $\mu_i$  is a correlated equilibrium so  $\kappa_i(A \times (0, 1]) = -\kappa_i(A \times [-1, 0))$ . Hence  $|\kappa_i|(g^{-1}(B)) = |\kappa_i|(A \times [-1, 0)) = |\kappa_i|(A \times (0, 1]) = |\kappa_i|(B)$ . Again, the proof is the same for  $B$  contained in other quadrants, so  $g$  is  $|\kappa_i|$ -measure preserving.

For the second-to-last step we prove that  $g$  is ergodic with respect to  $|\kappa|$ . Suppose  $B \subseteq I \times I$  is such that  $|\kappa|(g^{-1}(B) \triangle B) = 0$ . Let  $Q_i$  be the intersection of  $B$  with the  $i^{\text{th}}$  quadrant. Then  $|\kappa|(g^{-1}(Q_{i+1}) \triangle Q_i) = 0$ , so  $|\kappa|(g^{-4}(Q_1) \triangle Q_1) = 0$ . Let  $A = j_1^{-1}(Q_1)$ . Then  $|\kappa|(g^{-4}(Q_1) \triangle Q_1) = \nu_1((f_4 \circ f_3 \circ f_2 \circ f_1)^{-1}(A) \triangle A) = 0$ . By assumption the map  $f_4 \circ f_3 \circ f_2 \circ f_1$  is ergodic, so  $\nu_1(A) = 0$  or  $\nu_1(A) = \nu_1((0, 1]) = |\kappa|((0, 1] \times (0, 1])$ . Therefore  $|\kappa|(Q_1) = \nu_1(A) = 0$  or  $|\kappa|(Q_1) = |\kappa|((0, 1] \times (0, 1])$ . In either case since  $g$  is  $|\kappa|$ -measure preserving we get  $|\kappa|(Q_i) = |\kappa|(Q_1)$  for all  $i$ . Therefore  $|\kappa|(B) = 0$  or  $|\kappa|(B) = |\kappa|(I \times I)$ , so  $g$  is ergodic with respect to  $|\kappa|$ .

Normalizing  $|\kappa|$  and  $|\kappa_i|$  to be probability measures, we can apply Corollary 4.11 to obtain  $|\kappa_i| = \frac{|\kappa_i|(I \times I)}{|\kappa|(I \times I)} |\kappa|$ . By assumption the set on which  $|xy|$  is zero has  $\mu$  measure zero. Therefore

$$d\mu_i = \frac{1}{|xy|} d|\kappa_i| = \frac{|\kappa_i|(I \times I)}{|\kappa|(I \times I)} \frac{1}{|xy|} d|\kappa| = \frac{|\kappa_i|(I \times I)}{|\kappa|(I \times I)} d\mu,$$

so  $\mu_i = \frac{|\kappa_i|(I \times I)}{|\kappa|(I \times I)} \mu$  and  $\mu$  is extreme. □

Above we have constructed  $\mu$  and  $g$  so that  $g$  maps the quadrants counter-clockwise – quadrant 1 to quadrant 2, etc. However, the same argument would go through if  $g$  mapped the quadrants clockwise.

To view Proposition 4.5 as a special case of Proposition 4.12, let each  $\nu_i$  be a uniform probability measure over a finite subset of  $(0, 1]$ . The function  $g$  is defined by  $g(x_i, y_i) = (x_{i+1}, y_{i+1})$  and the  $f_i$  are defined to be compatible with this. The map  $f_4 \circ f_3 \circ f_2 \circ f_1$  is a permutation on the support of  $\nu_1$ , which is precisely the positive values of  $x_i$ . By construction this permutation consists of a single cycle, hence it is ergodic.

*Example 2.3 (cont'd).* We can combine Example 4.9 and Proposition 4.12 to exhibit extreme points of the set of correlated equilibria for this game which are not finitely supported. Let  $0 < a < b < 1$ . Let  $\nu_i$  be Lebesgue measure on  $[a, b)$  for all  $i$ . Fix  $\alpha$  such that  $\frac{\alpha}{b-a}$  is irrational. Define  $f_1 : [a, b) \rightarrow [a, b)$  by  $f(x) = (x - a + \alpha \bmod (b - a)) + a$ . This is just an affinely scaled version of Example 4.9 so  $f_1$  is  $\nu_i$ -measure preserving and ergodic. Define  $f_1$  on  $(0, 1] \setminus [a, b)$  arbitrarily, because that is a set of measure zero. Let  $f_2, f_3, f_4 : (0, 1] \rightarrow (0, 1]$  be the identity. These data satisfy all the assumptions of Proposition 4.12. In particular, since  $0 < a < b < 1$ ,  $xy$  is bounded away from zero on the support of  $|\kappa|$ . Therefore  $\frac{1}{|xy|} \in \mathcal{L}^1(|\kappa|)$ . Since  $\nu_i$  is not finitely supported,  $\mu$  is an extreme correlated equilibrium which is not finitely supported. The support of  $\mu$  is shown in Figure 3 with parameters  $a = 0.2$ ,  $b = 0.8$ , and  $\alpha = \frac{1}{\sqrt{5}}$ .

**Definition 4.13.** Given a compact Hausdorff space  $K$  we say that a set of measures  $\mathcal{M} \subseteq \Delta^*(K)$  is **describable by moments** if there exists an integer  $d$ , continuous maps




Figure 3: The support set of an extreme correlated equilibrium which is not finitely supported. Extremality of this equilibrium depends sensitively on the choices of endpoints for the line segments. In this case there are segments connecting:  $(0.2, -0.2)$  to  $(0.8, -0.8)$ ;  $(-0.2, -0.2)$  to  $(-0.8, -0.8)$ ;  $(-0.2, 0.2)$  to  $(-0.8, 0.8)$ ;  $(0.2, 0.2 + \frac{1}{\sqrt{5}})$  to  $(0.8 - \frac{1}{\sqrt{5}}, 0.8)$ ; and  $(0.8 - \frac{1}{\sqrt{5}}, 0.2)$  to  $(0.8, 0.2 + \frac{1}{\sqrt{5}})$ .

$g_1, \dots, g_d : K \rightarrow \mathbb{R}$ , and a set  $M \subseteq \mathbb{R}^d$  such that a measure  $\mu$  is in  $\mathcal{M}$  if and only if  $(\int g_1 d\mu, \dots, \int g_d d\mu) \in M$ .

The results of [12] show that the maximin and minimax strategy sets of a two-player zero-sum polynomial game can always be described by moments. Introducing a similar notion for  $n$ -tuples of moments, the set of Nash equilibria can always be described by moments in any polynomial game [22]. However, combining this example with the following proposition we see that the set of correlated equilibria of a polynomial game cannot in general be described by moments.

**Proposition 4.14.** *Let  $\mathcal{M} \subseteq \Delta^*(K)$  be a set of measures describable by moments. Then all extreme points of  $\mathcal{M}$  have finite support and this support is uniformly bounded by  $d$ , where  $d$  is the integer associated with the description of  $\mathcal{M}$  by moments.*

*Proof.* Suppose there exists a measure  $\mu \in \mathcal{M}$  which is extreme and supported on more than  $d$  points, so we can partition the domain of  $\mu$  into  $d+1$  sets  $B_1, \dots, B_{d+1}$  of positive measure. For  $c = (c_1, \dots, c_{d+1}) \in \mathbb{R}^{d+1}$ , define  $\mu_c = \sum_{i=1}^{d+1} c_i \mu|_{B_i}$ . The map  $c \mapsto \mu_c$  is injective. Define

$$K = \{c \in \mathbb{R}^{d+1} \mid \mu_c \text{ has the same moments as } \mu\},$$

so  $(1, 1, \dots, 1) \in K$ . By linearity of integration and the assumption that the  $B_i$  have positive measure,  $K$  is a convex polytope. By Carathéodory's theorem [2], the extreme points of  $K$  each have at most  $d$  nonzero entries. Thus  $(1, 1, \dots, 1)$  is not an extreme point of  $K$ , so we



can write  $(1, 1, \dots, 1) = \lambda c + (1 - \lambda)c'$  for  $0 < \lambda < 1$  and  $c, c' \neq (1, 1, \dots, 1)$ . Therefore  $\mu = \mu_{(1,1,\dots,1)} = \lambda\mu_c + (1 - \lambda)\mu_{c'}$  is not extreme.  $\square$

## 5 Future work

These results leave several open questions. If we define a moment map to be any map of the form  $\pi \mapsto (\int f_1 d\pi, \int f_2 d\pi, \dots, \int f_k d\pi)$  for continuous  $f_i$ , then we have shown that the set of correlated equilibria is not the inverse image of any set under any moment map. On the other hand, since moment maps are linear and weak\* continuous, we know that the image of the set of correlated equilibria under any moment map is convex and compact. Supposing the utilities and the  $f_i$  are polynomials, is there anything more we can say about this image? In particular, is it semialgebraic (i.e., describable in terms of finitely many polynomial inequalities)? If so, can we compute these inequalities efficiently for given utilities?

Another important question is: how typical is the complexity of extreme correlated equilibria observed above? What is the ratio of extreme correlated equilibria to extreme Nash equilibria “on average”? Are there examples where this ratio is higher than in our construction above?

## Acknowledgements

The first author would like to thank Prof. Cesar E. Silva for many discussions about ergodic theory, and in particular for the simple proof of Corollary 4.11 using Birkhoff’s ergodic theorem.

## References

- [1] R. J. Aumann. Subjectivity and correlation in randomized strategies. *Journal of Mathematical Economics*, 1(1):67 – 96, 1974.
- [2] D. P. Bertsekas, A. Nedić, and A. E. Ozdaglar. *Convex Analysis and Optimization*. Athena Scientific, Belmont, MA, 2003.
- [3] M. Cripps. Extreme correlated and Nash equilibria in two-person games. <http://www.olin.wustl.edu/faculty/cripps/CES2.DVI>, November 1995.
- [4] R. Datta. Universality of Nash equilibrium. *Mathematics of Operations Research*, 28(3):424 – 432, August 2003.
- [5] M. Dresher and S. Karlin. Solutions of convex games as fixed points. In H. W. Kuhn and A. W. Tucker, editors, *Contributions to the Theory of Games II*, number 28 in Annals of Mathematics Studies, pages 75 – 86. Princeton University Press, Princeton, NJ, 1953.

- [6] M. Dresher, S. Karlin, and L. S. Shapley. Polynomial games. In H. W. Kuhn and A. W. Tucker, editors, *Contributions to the Theory of Games I*, number 24 in Annals of Mathematics Studies, pages 161 – 180. Princeton University Press, Princeton, NJ, 1950.
- [7] F. S. Evangelista and T. E. S. Raghavan. A note on correlated equilibrium. *International Journal of Game Theory*, 25(1):35 – 41, March 1996.
- [8] D. Fudenberg and J. Tirole. *Game Theory*. MIT Press, Cambridge, MA, 1991.
- [9] F. Germano and G. Lugosi. Existence of sparsely supported correlated equilibria. *Economic Theory*, 32(3):575 – 578, September 2007.
- [10] I. L. Glicksberg. A further generalization of the Kakutani fixed point theorem, with application to Nash equilibrium points. *Proceedings of the American Mathematical Society*, 3(1):170 – 174, February 1952.
- [11] S. Hart and D. Schmeidler. Existence of correlated equilibria. *Mathematics of Operations Research*, 14(1), February 1989.
- [12] S. Karlin. *Mathematical Methods and Theory in Games, Programming, and Economics*, volume 2: Theory of Infinite Games. Addison-Wesley, Reading, MA, 1959.
- [13] S. Karlin and L. S. Shapley. *Geometry of Moment Spaces*. American Mathematical Society, Providence, RI, 1953.
- [14] R. J. Lipton, E. Markakis, and A. Mehta. Playing large games using simple strategies. In *Proceedings of the 4th ACM Conference on Electronic Commerce*, pages 36 – 41, New York, NY, 2003. ACM Press.
- [15] R. B. Myerson. *Game Theory: Analysis of Conflict*. Harvard University Press, Cambridge, MA, 1991.
- [16] J. F. Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286 – 295, September 1951.
- [17] R. Nau, S. G. Canovas, and P. Hansen. On the geometry of Nash equilibria and correlated equilibria. *International Journal of Game Theory*, 32:443 – 453, 2003.
- [18] W. Rudin. *Real & Complex Analysis*. WCB / McGraw-Hill, New York, 1987.
- [19] W. Rudin. *Functional Analysis*. McGraw-Hill, New York, 1991.
- [20] C. E. Silva. *Invitation to Ergodic Theory*. American Mathematical Society, Providence, RI, 2007.
- [21] N. D. Stein. Characterization and computation of equilibria in infinite games. Master’s thesis, Massachusetts Institute of Technology, May 2007.

- [22] N. D. Stein, A. Ozdaglar, and P. A. Parrilo. Separable and low-rank continuous games. *International Journal of Game Theory*, 37(4):475 – 504, December 2008.
- [23] N. D. Stein, P. A. Parrilo, and A. Ozdaglar. Correlated equilibria in continuous games: Characterization and computation. *Games and Economic Behavior*, to appear.